



Genetic Fact Sheets for Parents

Fatty Acid Oxidation Disorders

Screening, Technology, and Research in Genetics is a multi-state project to improve information about the financial, ethical, legal, and social issues surrounding expanded newborn screening and genetic testing – <http://www.newbornscreening.info>

Disorder Name: Short chain 3-hydroxyacyl-CoA dehydrogenase deficiency

Acronym: M/SCHADD

- What is M/SCHADD?
- What causes M/SCHADD?
- If M/SCHADD is not treated, what problems occur?
- What is the treatment for M/SCHADD?
- What happens when M/SCHADD is treated?
- What causes the M/SCHAD enzyme to be absent or not working correctly?
- How is M/SCHADD inherited?
- Is there genetic testing available?
- What other testing is available?
- Can you test during pregnancy?
- Can other members of the family have M/SCHADD or be carriers?
- Can other family members be tested?
- How many people have M/SCHADD?
- Does M/SCHADD happen more often in a certain ethnic group?
- Does M/SCHADD go by any other names?
- Where can I find more information?

This fact sheet contains general information about M/SCHADD. Every child is different and some of this information may not apply to your child specifically. Very little is known about M/SCHADD and, at present, there is no standard treatment plan. Certain treatments may be recommended for some children but not others. Children with M/SCHADD should be followed by a metabolic doctor in addition to their primary doctor.

What is M/SCHADD?

M/SCHADD stands for “medium/short chain 3-hydroxyacyl CoA dehydrogenase deficiency”. This disorder is also known as “3-hydroxyacyl CoA dehydrogenase deficiency”. It is one type of [fatty acid oxidation disorder](#). Some people with M/SCHADD have problems using fat for energy. However, most babies with newborn screening results showing M/SCHADD never have symptoms.

Fatty Acid Oxidation Disorders:

Fatty acid oxidation disorders (FAODs) are a group of rare inherited conditions. They are caused by enzymes that do not work properly.

A number of enzymes are needed to break down fats in the body (a process called fatty acid oxidation). Problems with any of these enzymes can cause a fatty acid oxidation disorder. People with FAODs cannot properly break down fat from either the food they eat or from fat stored in their bodies.


The symptoms and treatment vary between different FAODs. They can also vary from person to person with the same FAOD. See the fact sheets for each specific FAOD.

FAODs are inherited in an autosomal recessive manner and affect both males and females.

What causes M/SCHADD?

M/SCHADD occurs when an enzyme, called “short chain 3-hydroxyacyl CoA dehydrogenase” (SCHAD), is either missing or not working properly. This enzyme’s job is to help change certain fats in the food we eat into energy for the body. It also helps us use fat already stored in the body.

Short Chain 3-Hydroxyacyl-CoA Dehydrogenase Deficiency SCHADD



Energy from fat keeps us going whenever our bodies run low of their main source of energy, a type of sugar called glucose. Our bodies rely on fat when we don't eat for a stretch of time – like when we miss a meal or when we sleep.

Some people with M/SCHADD cannot break down fat for energy. Most babies found to have M/SCHADD on newborn screening do not seem to have this problem and do not develop symptoms.

If M/SCHADD is not treated, what problems occur?

M/SCHADD is highly variable and not well understood. Things that cause stress, such as lack of sleep, lack of food, illness or infection are thought to trigger episodes of illness called metabolic crises in some children with M/SCHADD but not in others.

Some of the first symptoms of a metabolic crisis are:

- extreme sleepiness
- behavior changes
- irritable mood

- poor appetite

Other symptoms then follow:

- fever
- diarrhea
- vomiting
- low blood sugar, called hypoglycemia

If a metabolic crisis is not treated, a child with M/SCHADD can develop:

- breathing problems
- seizures
- swelling of the brain
- coma, sometimes leading to death

Other effects of M/SCHADD can include:

- irregular heart beat and other heart problems
- enlarged heart
- liver problems
- muscle problems
- high levels of insulin in the blood in some babies

Symptoms of a metabolic crisis often happen after having nothing to eat for more than a few hours. Symptoms are also more likely when a person with M/SCHADD gets sick or has an infection.

What is the treatment for M/SCHADD?

Your baby's primary doctor may work with a metabolic doctor to care for your child. Your doctor may also suggest that you meet with a dietician familiar with M/SCHADD.

Certain treatments may be advised for some children but not others. When necessary, treatment is usually needed throughout life. The following are treatments that may be recommended for some children with M/SCHADD:

1. Avoid going a long time without food

Some infants and young children with M/SCHADD may need to eat frequently to prevent a metabolic crisis. Your metabolic doctor will tell you how often your child needs to be fed. In general, it is often suggested that infants that need to avoid fasting be fed every four to six hours. Some babies may need to eat even more frequently than this. It is important that these infants be fed during the night. If needed, your metabolic doctor and dietician will give you an appropriate feeding plan for your infant. Your doctor may also give you a 'sick day' plan tailored to your child's needs for you to follow during illnesses or other times when your child will not eat.

2. Diet

Sometimes a low fat, high carbohydrate food plan is recommended. Carbohydrates give the body many types of sugar that can be used as energy. In fact, for children needing this treatment, most food in the diet should be carbohydrates (bread, pasta, fruit, vegetables, etc.) and protein (lean meat and low-fat dairy foods). Any diet changes should be made under the guidance of a dietician familiar with M/SCHADD.

Ask your doctor whether your child needs to have any changes in his or her diet.

3. L-carnitine supplements and other medications

Some children may be helped by taking L-carnitine. This is a safe and natural substance that helps the body create energy. It also helps the body get rid of harmful wastes. Your doctor will decide whether your child needs L-carnitine. Unless you are advised otherwise, use only L-carnitine prescribed by your doctor.

Babies with high insulin levels may need medication. Your doctor will let you know if your child needs to take medication for this.

Do not use any medications or supplements without checking with your doctor.

4. Call your doctor at the start of any illness

If your child has shown previous symptoms of M/SCHADD, always call your health care provider when he or she has any of the following:

- poor appetite
- low energy or excessive sleepiness
- vomiting
- diarrhea
- an infection
- a fever
- persistent muscle pain or weakness

Children who have had symptoms of M/SCHADD need to eat extra starchy food and drink more fluids during any illness - even if they may not feel hungry - or they could develop a metabolic crisis. Children who are sick often don't want to eat. If they won't or can't eat, children with symptoms of M/SCHADD may need to be treated in the hospital to prevent problems.

Ask your metabolic doctor if you should carry a special travel letter with medical instructions for your child's care.

What happens when M/SCHADD is treated?

It is not known how effective treatment is in preventing health problems. It is hoped that with prompt and careful treatment children with M/SCHADD will be able to live healthy lives with typical growth and development.

The goal of treatment is to prevent death and serious health problems. However, children who have repeated episodes of hypoglycemia or metabolic crisis may develop brain damage. This can result in learning disabilities, intellectual disabilities or other lifelong effects.

What causes the SCHAD enzyme to be absent or not working correctly?

Genes tell the body to make various enzymes. People with M/SCHADD have a pair of genes that do not work correctly. Because of the changes in this pair of genes, the SCHAD enzyme either does not work properly or is not made at all.


How is M/SCHADD inherited?

M/SCHADD is inherited in an autosomal recessive manner. It affects both boys and girls equally.

Everyone has a pair of genes that make the SCHAD enzyme. In children with M/SCHADD, neither of these genes works correctly. These children inherit one non-working gene for the condition from each parent.

Parents of children with M/SCHADD rarely have the disorder. Instead, each parent has a single non-working gene for M/SCHADD. They are called carriers. Carriers do not have M/SCHADD because the other gene of this pair is working correctly.

When both parents are carriers, there is a 25% chance in each pregnancy for the child to have M/SCHADD. There is a 50% chance for the child to be a carrier, just like the parents. And, there is a 25% chance for the child to have two working genes.



Genetic counseling is available to families who have children with M/SCHADD. Genetic counselors can answer your questions about how M/SCHADD is inherited, choices during future pregnancies, and how to test other family members. Ask your doctor about a referral to a genetic counselor.

Is genetic testing available?

Genetic testing for M/SCHADD may be available. Genetic testing, also called DNA testing, look for changes in the pair of genes that cause M/SCHADD. Ask your metabolic doctor or genetic counselor about genetic testing for M/SCHADD.

DNA testing is not necessary to diagnose your child. It can be helpful for carrier testing or prenatal diagnosis, discussed below.

What other testing is available?

M/SCHADD can confirmed by a special test called a “fatty acid oxidation probe” using a skin sample. Talk to your doctor or your genetic counselor if you have questions about testing for M/SCHADD.

Can you test during pregnancy?

It may be possible to test for M/SCHADD during pregnancy either by DNA testing or through a special test called a “fatty acid oxidation probe” using cells from the fetus. The sample needed for these tests is obtained by either CVS or amniocentesis.

Parents can either choose to having testing during pregnancy or wait until birth to have the baby tested. A genetic counselor can talk to you about your choices and answer other questions you may have about prenatal testing or testing your baby after birth.

Can other members of the family have M/SCHADD or be carriers?

Having M/SCHADD

The brothers and sisters of a baby with M/SCHADD have a small chance of having this condition, even if they haven't had symptoms. Finding out whether other children in the family have M/SCHADD may be important because early treatment may prevent serious health problems. Talk to your doctor or genetic counselor about testing your other children for M/SCHADD.

M/SCHADD Carriers

Brothers and sisters who do not have M/SCHADD still have a chance to be carriers like their parents. Except in special cases, carrier testing should only be done in people over 18 years of age.

Each of the parents' brothers and sisters has a 50% chance to be a M/SCHADD carrier. It is important for other family members to be told that they could be carriers. There is a very small chance they are also at risk to have children with M/SCHADD.

Some states do not offer newborn screening for M/SCHADD. However, expanded newborn screening through private labs is available for babies born in states that do not screen for M/SCHADD. To learn more about expanded newborn screening, see [How to obtain MS/MS](#).

When both parents are M/SCHADD carriers, newborn screening results are not sufficient to rule out M/SCHADD in a newborn baby. In this case, special diagnostic testing should be done in addition to newborn screening.

During pregnancy, women carrying fetuses with M/SCHADD may be at increased risk to develop serious medical problems. Some women carrying fetuses with Fatty Acid Oxidation Disorders have developed:

- excessive vomiting
- abdominal pain
- high blood pressure
- jaundice
- abnormal fat storage in the liver
- severe bleeding

All women with a family history of M/SCHADD should share this information with their obstetricians and other health care providers before and during any future pregnancies. Knowing about these risks allows better medical care and early treatment if needed.

Can other family members be tested?

Diagnostic testing

Brothers and sisters can be tested for M/SCHADD using a skin sample. DNA testing using a blood sample may be possible, too.

Carrier Testing

Carrier testing for M/SCHADD may be available. Ask your metabolic doctor or genetic counseling whether carrier testing is possible.

How many people have M/SCHADD?

M/SCHADD is very rare. The actual incidence is unknown.

Does M/SCHADD happen more frequently in a certain ethnic group?

M/SCHADD does not happen more often in any specific race, ethnic group, geographical area or country.

Does M/SCHADD go by any other names?

M/SCHADD is sometimes also called:

- SCHAD deficiency
- SCHADD
- HADHSC deficiency
- HADH deficiency
- HAD deficiency
- L-3-alpha-hydroxyacyl-CoA dehydrogenase, short chain, deficiency

Where can I find more information?

Fatty Oxidation Disorders (FOD) Family Support Group

<http://www.fodsupport.org>

Organic Acidemia Association

<http://www.oaanews.org>

United Mitochondrial Disease Foundation

<http://www.umdf.org>

Children Living with Inherited Metabolic Diseases (CLIMB)

<http://www.climb.org/uk>

Baby's First Test

<http://www.babysfirsttest.org>

<u>Document Info</u>	Created by:	www.newbornscreening.info
	Reviewed by:	HI, CA, OR, and WA metabolic specialists
	Review date:	July 13, 2013 April 2, 2011 September 31, 2007
	Updated on:	February 28, 2016

DISCLAIMER:

THIS INFORMATION DOES NOT PROVIDE MEDICAL ADVICE. All content ("Content"), including text, graphics, images and information are for general informational purposes only. You are encouraged to confer with your doctor or other health care professional with regard to information contained on this information sheet. After reading this information sheet, you are encouraged to review the information carefully with your doctor or other healthcare provider. The Content is not intended to be a substitute for professional medical advice, diagnosis or treatment. NEVER DISREGARD PROFESSIONAL MEDICAL ADVICE, OR DELAY IN SEEKING IT, BECAUSE OF SOMETHING YOU HAVE READ ON THIS INFORMATION SHEET.



This project is supported by a grant from the Maternal and Child Health Bureau, Health Resources and Service Administration, Genetic Services Branch, MCH Project #:1H46 MC 00189-03 <http://mchb.hrsa.gov>

Short chain 3-hydroxyacyl-CoA dehydrogenase deficiency

Created by www.newbornscreening.info

10

Review Date: 2/28/2016

